

AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-305-671-0200.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Drug Requested: Palforzia® [Peanut (Arachis hypogaea) Allergen Powder-dnfp] (**Pharmacy**)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member AvMed #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight: _____ Date: _____

<u>Medication</u>	<u>Quantity Limit</u>
Palforzia Initial Dose Escalation Kit	1 kit per 365 days
Palforzia Up-Dosing Kits (Levels 1-11)	1 kit per 365 days
Palforzia 300 mg sachets	1 sachet per day

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 12 months

- Member must have diagnosis of peanut allergy
- Member must be at least 4-17 years of age at initiation of therapy
- Prescribed by or in consultation with an Allergist or Immunologist

(Continued on next page)

- Provider has submitted documentation to confirm diagnosis of peanut allergy via **ONE** of the following:
 - Member has a diagnosis and clinical history of peanut allergy as documented by **BOTH** of the following (**must submit labs and skin prick test results for documentation**):
 - A serum peanut-specific IgE level of ≥ 0.35 kUA/L
 - A mean wheal diameter that is at least 3 mm larger than the negative control on skin prick test for peanut
 - In the absence of positive clinician supervised food challenge, peanut allergy is confirmed by the **BOTH** of the following:
 - Positive skin prick test to peanut ≥ 8 mm compared to control, unless skin testing is contraindicated
 - Serum IgE to peanut ≥ 14 kUA/L
- Palforzia will be used in conjunction with a peanut-avoidance diet Member must be prescribed injectable epinephrine (**verified by chart notes or pharmacy paid claims**)
- Member and/or caregiver has been instructed and trained on the appropriate use of injectable epinephrine
- Health care provider, health care setting, and member **MUST** be enrolled in the Palforzia REMS program
- Request for Palforzia may **NOT** be approved if member has **ANY** of the following:
 - Severe or poorly controlled asthma
 - History of eosinophilic esophagitis or other eosinophilic gastrointestinal disease
 - History of severe or life-threatening episodes of anaphylaxis or anaphylactic shock within the past 2 months
 - History of mast cell disorder (including mastocytosis), urticarial pigmentosa, hereditary or idiopathic angioedema or currently has paid claims for Berinert, Cinryze, Haegarda, Firazyr, Takhyzyro or Ruconest
 - Individual is in buildup phase of immunotherapy to another allergen (i.e. has not reached maintenance dosing)

Reauthorization: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Provider please note: a one-time reauthorization is required after initial 12 month approval

- Member must continue to tolerate the prescribed daily dose of Palforzia[®]
- Member is compliant with Palforzia[®] therapy (**verified by pharmacy paid claims**)
- Member has **NOT** experienced recurrent asthma exacerbations
- Member has **NOT** experienced any treatment-restricting adverse effects (e.g., repeated systemic allergic reaction and/or severe anaphylaxis)

(Continued on next page)

Dosing Tables

Table 1: Dosing Configuration for Initial Dose Escalation (Single Day Dose Escalation); supplied as a single card consisting of 5 blisters containing a total of 13 capsules

Dose Level	Total Dose	Dose Configuration
A	0.5 mg	One 0.5 mg capsule
B	1 mg	One 1 mg capsule
C	1.5 mg	One 0.5 mg capsule; One 1 mg capsule
D	3 mg	Three 1 mg capsules
E	6 mg	Six 1 mg capsules

Table 2: Daily Dosing Configuration for Up-Dosing

Dose Level	Total Daily Dose	Daily Dose Configuration	Dose Duration (weeks)
1	3 mg	Three 1 mg capsules	2
2	6 mg	Six 1 mg capsules	2
3	12mg	Two 1 mg capsules; One 10 mg capsule	2
4	20 mg	One 20 mg capsule	2
5	40 mg	Two 20 mg capsules	2
6	80 mg	Four 20 mg capsules	2
7	120 mg	One 20 mg capsule; One 100 mg capsule	2
8	160 mg	Three 20 mg capsules; One 100 mg capsule	2
9	200 mg	Two 100 mg capsules	2
10	240 mg	Two 20 mg capsules; Two 100 mg capsules	2
11	300 mg	One 300 mg sachet	2

Table 3: Daily Dosing Configuration for Maintenance

Dose Level	Total Daily Dose	Daily Dose Configuration
11	300 mg	One 300 mg sachet

Medication being provided by a Specialty Pharmacy - PropriumRx

Not all drugs may be covered under every Plan

If a drug is non-formulary on a Plan, documentation of medical necessity will be required.

Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.