

AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-305-671-0200.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Drug Requested: Rezdifra™ (resmetirom)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member AvMed #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight: _____ Date: _____

Recommended Dosing: <100 kg: 80 mg once daily. ≥100 kg: 100 mg once daily. Coadministration with moderate CYP2C8 inhibitors: Reduce dose to 80 mg daily for patients weighing ≥100 kg, or reduce dose to 60 mg daily for patients weighing <100 kg

Quantity Limits: One tablet daily (all strengths – 60, 80 & 100 mg)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 12 months

- Member is 18 years of age or older
- Medication is prescribed by or in consultation with a hepatologist or gastroenterologist
- Member has a diagnosis of nonalcoholic steatohepatitis or metabolic dysfunction-associated steatohepatitis (NASH/MASH)

(Continued on next page)

- ❑ Provider must submit **ONE** of the following
 - ❑ Biopsy results (completed within the last 6 months) documenting **BOTH** of the following:
 - ❑ Liver fibrosis stage F2 or F3
 - ❑ Non-alcoholic fatty liver disease activity score (NAS) of ≥ 4 with a score of > 1 in all the following: steatosis, ballooning, and lobular inflammation
 - ❑ Liver fibrosis stage F2 or F3 as determined by an elastography test, such as vibration-controlled transient elastography (i.e., FibroScan), magnetic resonance elastography (MRE), shear wave elastography; etc. (**must submit current test results**)
- ❑ In cases of indeterminate fibrosis stage (i.e., inconsistency between fibrosis stage and clinical presentation), a liver biopsy will be required to be submitted
- ❑ Member has three or more of the following metabolic risk factors that are managed according to standard of care (**verified by medical chart notes, lab test results and/or pharmacy claims**):
 - ❑ Central obesity
 - ❑ Hypertriglyceridemia
 - ❑ Reduced high-density lipoprotein cholesterol
 - ❑ Hypertension
 - ❑ Elevated fasting plasma glucose indicative of diabetes or pre-diabetes
- ❑ Current liver function (CMP) and CBC test results must be submitted
- ❑ Other causes of liver disease or hepatic steatosis have been ruled out (i.e., alcoholic steatohepatitis, acute fatty liver, autoimmune hepatitis, Hepatitis A, B or C, hemochromatosis, drug-induced liver disease)
- ❑ Member has adopted liver-protective lifestyle interventions such as optimizing weight loss, dietary changes, and exercise
- ❑ Member does **NOT** have significant alcohol consumption (alcohol consumption of more than 20 g per day for women and more than 30 g per day for men)
- ❑ Member does **NOT** have evidence of cirrhosis, hepatic decompensation, or hepatocellular carcinoma (**must submit documentation**)

Reauthorization: 6 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- ❑ Member continues to meet **ALL** initial authorization criteria
- ❑ Member has experienced **ONE** of the following as determined by an elastography test, such as vibration-controlled transient elastography (e.g., FibroScan), magnetic resonance elastography (MRE), shear wave elastography or biopsy:
 - ❑ MASH/NASH resolution **AND** no worsening of fibrosis
 - ❑ No worsening of MASH/NASH **AND** improvement in fibrosis by ≥ 1 stage

(Continued on next page)

Medication being provided by Specialty Pharmacy – Proprium Rx

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****