AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-305-671-0200</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

Drug Requested: Actemra® SQ (tocilizumab) (Pharmacy)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.				
Member Name:				
Member AvMed #:	Date of Birth:			
Prescriber Name:				
Prescriber Signature:	Date:			
Office Contact Name:				
Phone Number:	Fax Number:			
DEA OR NPI #:				
DRUG INFORMATION: Authorization may be delayed if incomplete.				
Drug Form/Strength:				
	Length of Therapy:			
Diagnosis:	ICD Code:			
Weight:	Date:			
\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	oncomitant therapy with more than one biologic mira, Rinvoq, Stelara) prescribed for the same or different al. Safety and efficacy of these combinations has NOT been			
	I that apply. All criteria must be met for approval. To ncluding lab results, diagnostics, and/or chart notes, must be			
□ Diagnosis: Moderate-to-Severe Rheumatoid Arthritis Dosing: SubQ: <100 kg - 162 mg once every other week; >100 kg - 162 mg once every week				
☐ Member has a diagnosis of moderate-to-s	severe rheumatoid arthritis			
☐ Prescribed by or in consultation with a R	heumatologist			

(Continued on next page)

		ember has tried and failed at least ONE of the following DMARD therapies for at least three (3) onths
		hydroxychloroquine
		leflunomide
		methotrexate
		sulfasalazine
	ember meets ONE of the following:	
		Member tried and failed, has a contraindication, or intolerance to ONE of the following PREFERRED adalimumab products:
		□ Humira®
		□ Cyltezo®
		□ Hyrimoz [®]
		*NOTE: Humira NDC's starting with 83457 are not approved, NDC's starting with 00074 (MFG: Abbvie) are preferred; Hyrimoz NDC's starting with 83457 are not approved, NDC's starting with 61314 (MFG: Sandoz) are preferred
		Member has been established on Actemra® for at least 90 days <u>AND</u> prescription claims history indicates <u>at least a 90-day supply of Actemra was dispensed within the past 130 days</u> (verified by chart notes or pharmacy paid claims)
	_	nosis: Juvenile Idiopathic Arthritis ng: SubQ: <30 kg – 162 mg once every 3 weeks; ≥30 kg – 162 mg once every 2 weeks
	osin	
D	osin Me	ig: SubQ: <30 kg − 162 mg once every 3 weeks; ≥30 kg − 162 mg once every 2 weeks
D	osin Me	rg: SubQ: $<30 \text{ kg} - 162 \text{ mg}$ once every 3 weeks; $\ge 30 \text{ kg} - 162 \text{ mg}$ once every 2 weeks ember is ≥ 2 years of age and has a diagnosis of $\underline{\mathbf{ONE}}$ of the following:
D	osin Me	ng: SubQ: <30 kg − 162 mg once every 3 weeks; ≥30 kg − 162 mg once every 2 weeks ember is ≥ 2 years of age and has a diagnosis of <u>ONE</u> of the following: Active polyarticular juvenile idiopathic arthritis (PJIA)
D	osin Me Pre	ng: SubQ: <30 kg − 162 mg once every 3 weeks; ≥30 kg − 162 mg once every 2 weeks ember is ≥ 2 years of age and has a diagnosis of <u>ONE</u> of the following: Active polyarticular juvenile idiopathic arthritis (PJIA) Active systemic juvenile idiopathic arthritis
D	osin Me Pre	ng: SubQ: <30 kg − 162 mg once every 3 weeks; ≥30 kg − 162 mg once every 2 weeks ember is ≥ 2 years of age and has a diagnosis of \underline{ONE} of the following: Active polyarticular juvenile idiopathic arthritis (PJIA) Active systemic juvenile idiopathic arthritis escribed by or in consultation with a Rheumatologist ember has tried and failed at least \underline{ONE} of the following DMARD therapies for at least \underline{three} (3)
D	osin Me Pre Me	ng: SubQ: <30 kg − 162 mg once every 3 weeks; ≥30 kg − 162 mg once every 2 weeks ember is ≥ 2 years of age and has a diagnosis of <u>ONE</u> of the following: Active polyarticular juvenile idiopathic arthritis (PJIA) Active systemic juvenile idiopathic arthritis escribed by or in consultation with a Rheumatologist ember has tried and failed at least <u>ONE</u> of the following DMARD therapies for at least three (3) onths
D	osin Me Pre Me	ng: SubQ: <30 kg − 162 mg once every 3 weeks; ≥30 kg − 162 mg once every 2 weeks ember is ≥ 2 years of age and has a diagnosis of <u>ONE</u> of the following: Active polyarticular juvenile idiopathic arthritis (PJIA) Active systemic juvenile idiopathic arthritis escribed by or in consultation with a Rheumatologist ember has tried and failed at least <u>ONE</u> of the following DMARD therapies for at least three (3) onths cyclosporine
D	Me D Pre Me mo	ng: SubQ: <30 kg − 162 mg once every 3 weeks; ≥30 kg − 162 mg once every 2 weeks ember is ≥ 2 years of age and has a diagnosis of <u>ONE</u> of the following: Active polyarticular juvenile idiopathic arthritis (PJIA) Active systemic juvenile idiopathic arthritis escribed by or in consultation with a Rheumatologist ember has tried and failed at least <u>ONE</u> of the following DMARD therapies for at least three (3) onths cyclosporine hydroxychloroquine
D	Me D Prec Me Me D D D D D D D D D D D D D D D D D	ember is ≥ 2 years of age and has a diagnosis of <u>ONE</u> of the following: Active polyarticular juvenile idiopathic arthritis (PJIA) Active systemic juvenile idiopathic arthritis escribed by or in consultation with a Rheumatologist ember has tried and failed at least <u>ONE</u> of the following DMARD therapies for at least three (3) onths cyclosporine hydroxychloroquine leflunomide
D	osin Me Pre Me mo	ember is ≥ 2 years of age and has a diagnosis of ONE of the following: Active polyarticular juvenile idiopathic arthritis (PJIA) Active systemic juvenile idiopathic arthritis escribed by or in consultation with a Rheumatologist ember has tried and failed at least ONE of the following DMARD therapies for at least three (3) onths cyclosporine hydroxychloroquine leflunomide methotrexate
D	Me D Pre Me Me D D D D D D D D D D D D D D D D	ember is ≥ 2 years of age and has a diagnosis of ONE of the following: Active polyarticular juvenile idiopathic arthritis (PJIA) Active systemic juvenile idiopathic arthritis escribed by or in consultation with a Rheumatologist ember has tried and failed at least ONE of the following DMARD therapies for at least three (3) onths cyclosporine hydroxychloroquine leflunomide methotrexate Non-steroidal anti-inflammatory drugs (NSAIDs)

(Continued on next page)

	For members with a diagnosis of PJIA only , member must meet ONE of the following:		
		Member tried and failed, has a contraindication, or intolerance to <u>ONE</u> of the following <u>PREFERRED</u> adalimumab products:	
		☐ Humira [®]	
		□ Cyltezo [®]	
		□ Hyrimoz [®]	
		*NOTE: Humira NDC's starting with 83457 are not approved, NDC's starting with 00074 (MFG: Abbvie) are preferred; Hyrimoz NDC's starting with 83457 are not approved, NDC's starting with 61314 (MFG: Sandoz) are preferred	
		Member has been established on Actemra® for at least 90 days AND prescription claims history indicates at least a 90-day supply of Actemra was dispensed within the past 130 days (verified by chart notes or pharmacy paid claims)	
Medication being provided by Specialty Pharmacy – Proprium Rx			

**Use of samples to initiate therapy does not meet step edit/preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *