## AvMed

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-305-671-0200</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

<b>Drug Requested:</b> (Select drug	below)		
□ <b>Libervant</b> <sup>™</sup> (diazepam) buccal film	□ Nayzilam <sup>®</sup> (midazolam nasal spray)		
MEMBER & PRESCRIBI	ER INFORMATION: Authorizat	ion may be delayed if incomplete.	
Member Name:			
	Date of Birth:		
Prescriber Name:			
Prescriber Signature:			
Office Contact Name:			
	Fax Number:		
NPI #:			
	Authorization may be delayed if incom		
Drug Name/Form/Strength:			
Dosing Schedule:		Therapy:	
	ICD Code, if applicable:		
Weight (if applicable):	Date weight obtained:		

## **Recommended Dosing:**

- **Libervant**<sup>™</sup> (diazepam) buccal films: Administer 5 mg to 15 mg orally, dependent on the patient's weight. A second dose, when required, may be administered at least 4 hours after the first dose. More than 2 doses of Libervant should not be used to treat a single episode. More than one episode should not be treated with Libervant every five days or more than five episodes per month.
- Nayzilam® (midazolam) nasal spray: Administer one spray (5 mg dose) into one nostril. Optional second dose after 10 minutes. Maximum dose: 2 doses per episode, 1 episode every 3 days, 5 episodes per month.
- Valtoco® (diazepam) nasal spray: Initial Dose: 5 mg and 10 mg doses are administered as a single spray intranasally into one nostril. Administration of 15 mg and 20 mg doses requires two nasal spray devices, one spray (7.5 mg or 10 mg) into each nostril. A second dose, when required, maybe administered at least 4 hours after the initial dose. Maximum dose: 2 doses per episode, 1 episode every 5 days, 5 episodes per month.
- Quantity Limit for Libervant<sup>™</sup>, Nayzilam<sup>®</sup> and Valtoco<sup>®</sup>: 10 spray units/films per 30 days

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

A 41	•	4 ·	$\sim$	•
	horiza	ition (	'P11	teria
$\Delta u u$		LLIVII		LUI IA

Member must meet <b>ONE</b> of the following age requirements:
☐ If requesting <b>Libervant</b> <sup>™</sup> , member must be between 2 to 5 years of age
☐ If requesting Nayzilam®, member must be 12 years of age or older
☐ If requesting Valtoco®, member must be 6 years of age or older
<u>AND</u>
Prescribing physician is a neurologist or has consulted with a neurologist
<u>AND</u>
Member has a diagnosis of epilepsy
<u>AND</u>
Member will be using requested medication for the acute treatment of intermittent episodes of frequent seizure activity (i.e., seizure clusters, acute repetitive seizures) which are distinct from the member's usual seizure pattern with epilepsy (chart notes must be submitted for documentation of seizure activity)
<u>AND</u>
Member is currently receiving maintenance antiepileptic medication(s) e.g., lamotrigine, levetiracetam topiramate, oxcarbazepine (verified by chart notes and/or pharmacy paid claims)
AND
Prescriber agrees to assess the member before prescribing concomitant opioid therapy to limit opioid dosages and durations to the minimum required
<u>AND</u>
Dose does not exceed the FDA-approved maximum dose
<u>AND</u>
Libervant <sup>™</sup> , Nayzilam <sup>®</sup> and Valtoco <sup>®</sup> will <u>NOT</u> be used concomitantly

Exclusion Criteria: Patients with known hypersensitivity to midazolam and acute narrow-angle glaucoma

\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. \*\*

\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. \*