AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-305-671-0200</u>. No additional phone calls will be necessary if all information <u>(including phone and fax #s)</u> on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

Drug Requested: Skyrizi® SQ & IV (risankizumab) For CD & UC (Pharmacy) (Preferred)

MEMBER & PRESCRIBE	R INFORMATION: Authorization may be delayed if incomplete.
Member Name:	
Member AvMed #:	Date of Birth:
Prescriber Name:	
	Date:
Office Contact Name:	
Phone Number:	
NPI #:	
DRUG INFORMATION: A	Authorization may be delayed if incomplete.
Drug Name/Form/Strength:	
Dosing Schedule:	Length of Therapy:
Diagnosis:	ICD Code, if applicable:
Weight (if applicable):	Date weight obtained:
	ng dose for treatment of Crohn's disease & Ulcerative colitis can only be EFIT . NDC: 00074-5015-01; J2327; 600 mg = 600 billable units, 1200 mg =
Adult Dosing:	
☐ Induction IV: NDC: 00074-50	15-01 – Skyrizi IV 600 mg/10 mL – J2327
□ Crohn's disease -600 mg and 8; $600 \text{ mg} = 600 \text{ billable}$	administered by IV infusion over a period of at least one hour at week 0, 4 e units per dose
☐ Ulcerative colitis – 1200 m 8; 1200 mg = 1200 billable	g administered by IV infusion a period of at least two hours at week 0, 4 and units per dose
00074-1065-01 – Skyrizi SQ 18	074-1069-01/00074-1070-01 — Skyrizi SQ 360 mg/ 2.4 mL cartridge; NDC 80 mg/ 1.2 mL cartridge. *Use the lowest effective dosage to maintain
therapeutic response*.	k 12, and every 8 weeks thereafter
<u>-</u>	k 12. and every 8 weeks thereafter

(Continued on next page)

(Continued on next page)

indications to be experimental and investigational. Satestablished and will NOT be permitted.	fety and efficacy of these combinations has NOT been
• Will the member be discontinuing a previously pre-	escribed biologic if approved for requested medication? — Yes OR — No
• If yes, please list the medication that will be discorapproval along with the corresponding effective data	ntinued and the medication that will be initiated upon ate.
Medication to be discontinued:	Effective date:
Medication to be initiated:	Effective date:
CLINICAL CRITERIA: Check below all that support each line checked, all documentation, include provided or request may be denied.	t apply. All criteria must be met for approval. To ing lab results, diagnostics, and/or chart notes, must be
☐ Maintenance Dose – 180 mg or 360 mg a week 12, and every 8 weeks thereafter	administered by subcutaneous injection at
<u>Authorization Criteria</u> : To be reviewed for	approval under the pharmacy benefit
☐ Member has <u>ONE</u> of the following diagnoses	
☐ Moderate-to-severe active Crohn's disease	e
☐ Moderate-to-severe active Ulcerative colit	is
 Prescribed by or in consultation with a Gastro 	enterologist
☐ Member meets <u>ONE</u> of the following:	
Member has tried and failed budesonide or	high dose steroids (40-60 mg prednisone)
Member has tried and failed at least <u>ONE</u> of <u>months</u>	of the following DMARD therapies for at least three (3)
 5-aminosalicylates (balsalazide, olsalaz 	zine, sulfasalazine)
☐ oral mesalamine (Apriso, Asacol/HD, I	Delzicol, Lialda, Pentasa)
☐ Induction Dose (If required) — One time to receive up to three (3) IV infusion do	e approval for duration of 2 months, member ses
<u>Authorization Criteria</u> : To be reviewed for	r one-time approval under the medical benefit
☐ Medication will be used as induction therapy	

(Continued on next page)

NOTE: The Health Plan considers the use of concomitant therapy with more than one biologic

immunomodulator (e.g., Dupixent, Entyvio, Humira, Rinvoq, Stelara) prescribed for the same or different

PA Skyrizi for CD & UC (Pharmacy)(AvMed) (Continued on next page)

	Me	edication being provided by:
		Location/site of drug administration:
		NPI or DEA # of administering location:
	Me	ember to receive FDA approved loading dose for ONE of the following indications:
		Crohn's disease $-600~\mathrm{mg}$ administered by IV infusion over a period of at least one hour at week 0, 4 and 8
		Ulcerative colitis -1200 mg administered by IV infusion a period of at least two hours at week $0,4$ and 8
Med	lica	tion being provided by a Specialty Pharmacy – Proprium Rx

**Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *