AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-305-671-0200</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

Preferred Adalimumab Products (Pharmacy)

Drug Requested: select one drug below

COMMERCIA	L/FAMIS PREFERREDS
□ Cyltezo® (adalimumab-adbm) □ Hum	ira® (adalimumab)
SMALL GROUP/INDIVIDUAL	PRODUCT/EXCHANGE PREFERREDS
□ adalimumab-adbm	□ Simlandi [®] (adalimumab-ryvk)
NOTE: Humira NDC's starting with 83457 are Note Preferred	OT approved, NDC's starting with 00074 (MFG: Abbvie) are
MEMBER & PRESCRIBER INFORM	ATION: Authorization may be delayed if incomplete.
Member Name:	
Member AvMed #:	Date of Birth:
Prescriber Name:	
Prescriber Signature:	Date:
Office Contact Name:	
Phone Number:	Fax Number:
NPI #:	
DRUG INFORMATION: Authorization m	ay be delayed if incomplete.
Drug Name/Form/Strength:	
Dosing Schedule:	Length of Therapy:
Diagnosis:	ICD Code, if applicable:
Weight (if applicable):	Date weight obtained:
	omitant therapy with more than one biologic ra, Rinvoq, Stelara) prescribed for the same or different Safety and efficacy of these combinations has NOT been

(Continued on next page)

PA Adalimumab Products (AvMed) (Continued from previous page)

• W	ill tl	he member be discontinuing a previously	prescribed biologic if approved for requested medication?
			☐ Yes OR ☐ No
	•	please list the medication that will be disval along with the corresponding effective	scontinued and the medication that will be initiated upon the date.
M	ledic	cation to be discontinued:	Effective date:
M	ledic	cation to be initiated:	Effective date:
supp	ort e		nat apply. All criteria must be met for approval. To luding lab results, diagnostics, and/or chart notes, must be agnosis below that applies.
		gnosis: Moderate-to-Severe Rheung: SubQ: 40 mg every other week	matoid Arthritis
	Me	ember has a diagnosis of moderate-to-sev	vere rheumatoid arthritis
	Pro	escribed by or in consultation with a Rhe	eumatologist
		ember has tried and failed at least <u>ONE</u> o onths	of the following DMARD therapies for at least three (3)
		hydroxychloroquine	
		leflunomide	
		methotrexate	
		sulfasalazine	
	_	gnosis: Moderate-to-Severe Activ ng: SubQ: 40 mg every other week	e Polyarticular Juvenile Idiopathic Arthritis
	M	ember has a diagnosis of moderate-to-sev	rere active polyarticular juvenile idiopathic arthritis
	Pro	escribed by or in consultation with a Rhe	umatologist
	Me	ember is ≥ 2 years of age	
	<u>m</u>	onths	of the following DMARD therapies for at least three (3)
		cyclosporine	
	_	hydroxychloroquine	
	_	leflunomide	
		methotrexate	NGATE)
		non-steroidal anti-inflammatory drugs (NSAIDS)
		oral corticosteroids	
		sulfasalazine	
		tacrolimus	

(Continued on next page)

2

	iagnosis: Active Psoriatic Arthritis osing: SubQ: 40 mg every other week	
	Member has a diagnosis of active psoriatic arthri	tis
	Prescribed by or in consultation with a Rheumatol	logist
	Member has tried and failed at least ONE of the formonths	bllowing DMARD therapies for at least three (3)
	□ cyclosporine	
	□ leflunomide	
	□ methotrexate	
	□ sulfasalazine	
	iagnosis: Active Ankylosing Spondylitis osing: SubQ: 40 mg every other week.	
	Member has a diagnosis of active ankylosing spor	ndylitis
	Prescribed by or in consultation with a Rheumatol	logist
	Member tried and failed, has a contraindication, or	intolerance to <u>TWO</u> NSAIDs
Do	iagnosis: Moderate-to-Severe Hidradenitiosing: SubQ: Initial: 160 mg (given on day 1 or spweeks later (day 15). Maintenance: 40 mg every	lit and given over 2 consecutive days); then 80 mg
	Member is ≥ 12 years of age and has a diagnosis of	f moderate-to-severe hidradenitis suppurativa
	Prescribed by or in consultation with a Dermatolo	gist
	Member tried and failed a 90-day course of oral an or clindamycin, rifampin) for treatment of HS (wit	tibiotics (e.g., tetracycline, minocycline, doxycycline hin last 9 months)
	Name of Antibiotic & Date:	
Do	iagnosis: Moderate-to-Severe Chronic Plansing: SubQ: Initial: 80 mg as a single dose. Main tek after initial dose.	-
	Member has a diagnosis of moderate-to-severe chr	onic plaque psoriasis
	Prescribed by or in consultation with a Dermatolo	gist
	Member tried and failed at least <u>ONE</u> of either Pholeast <u>three (3) months</u> (check each tried below):	ototherapy or Alternative Systemic Therapy for at
	□ Phototherapy:	☐ Alternative Systemic Therapy:
	☐ UV Light Therapy	□ Oral Medications
	□ NB UV-B	□ acitretin
	□ PUVA	☐ methotrexate
		□ cyclosporine

(Continued on next page)

□ Diagnosis: Moderate-to-Severe Active Cr Dosing: SubQ: Initial: 160 mg (given on day 1 or 2 weeks later (day 15). Maintenance: 40 mg every	split and given over 2 consecutive days); then 80 mg
\square Member is ≥ 6 years of age and has a diagnosis of	of moderate-to-severe active Crohn's disease
☐ Prescribed by or in consultation with a Gastroer	terologist
☐ Member meets <u>ONE</u> of the following:	
Member has tried and failed budesonide or h	igh dose steroids (40-60 mg prednisone)
 Member has tried and failed at least <u>ONE</u> of months 	the following DMARD therapies for at least three (3)
□ 5-aminosalicylates (balsalazide, olsalazin	e. sulfasalazine)
□ oral mesalamine (Apriso, Asacol/HD, De	,
	7
□ Diagnosis: Moderate-to-Severe Ulcerative Dosing: SubQ: Initial: 160 mg (given on day 1 OF mg 2 weeks later (day 15). Maintenance: 40 mg e	R split and given over 2 consecutive days); then 80
\square Member is ≥ 5 years of age and has a diagnosis of	of moderate-to-severe ulcerative colitis
☐ Prescribed by or in consultation with a Gastroe r	terologist
☐ Member meets <u>ONE</u> of the following:	
 Member has tried and failed budesonide or h 	igh dose steroids (40-60 mg prednisone)
Member has tried and failed at least <u>ONE</u> of <u>months</u>	the following DMARD therapies for at least three (3)
5-aminosalicylates (balsalazide, olsalazin	e, sulfasalazine)
☐ oral mesalamine (Apriso, Asacol/HD, De	Izicol, Lialda, Pentasa)
□ Diagnosis: Uveitis (non-infectious intermo Dosing: SubQ: Initial: 80 mg as a single dose. Ma week after initial dose.	
☐ Member is ≥ 2 years of age and has a diagnosis of applies):	of Uveitis (check box below for diagnosis that
□ Chronic	☐ Treatment-refractory
□ Recurrent	☐ Vision-threatening disease
☐ Prescribed by or in consultation with an Ophtha	lmologist or Rheumatologist

(Continued on next page)

Not all drugs may be covered under every Plan

If a drug is non-formulary on a Plan, documentation of medical necessity will be required.

**Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *