

AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-305-671-0200.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Drug Requested: COLONY STIMULATING FACTORS

[Form to be completed **ONLY** if the member is self-administering]

Short-acting Granulocyte Colony-Stimulating Factors (G-CSFs)		
<input type="checkbox"/> Granix [®] (tbo-filgrastim)	<input type="checkbox"/> Neupogen [®] (filgrastim)	<input type="checkbox"/> Nivestym [™] (filgrastim-aafi)
<input type="checkbox"/> Releuko [®] (filgrastim-ayow)	<input type="checkbox"/> Zarxio [®] (filgrastim-sndz)	
Granulocyte-macrophage Colony-Stimulating Factor (GM-CSF)		
<input type="checkbox"/> Leukine [®] (sargramostim)		
Long-acting Granulocyte Colony-Stimulating Factors (G-CSFs)		
<input type="checkbox"/> Fulphila [™] (pegfilgrastim-jmdb)	<input type="checkbox"/> Ryzneuta [®] (efbemalenograstim alfa-vuxw)	
<input type="checkbox"/> Fylnetra [™] (pegfilgrastim-pbbk)	<input type="checkbox"/> Stimufend [®] (pegfilgrastim-fpgk)	
<input type="checkbox"/> Neulasta [®] (pegfilgrastim)	<input type="checkbox"/> Udenyca [®] (pegfilgrastim-cbqv)	
<input type="checkbox"/> Nyvepria [™] (pegfilgrastim-apgf)	<input type="checkbox"/> Ziextenzo [™] (pegfilgrastim-bmez)	
<input type="checkbox"/> Rolvedon [™] (eflapegrastim-xnst)		

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member AvMed #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code: _____

Weight: _____ Date: _____

(Continued on next page)

Maximum Daily Dose:

Fulphila 6 mg prefilled syringe: 1 syringe/14 days	Nyvepria 6 mg prefilled syringe: 1 syringe/14 days
Fylnetra 6 mg prefilled syringe: 1 syringe/14 days	Releuko 300 mcg vial: 3 vials/1 day
Granix 300 mcg prefilled syringe: 4 syringes/1 day	Releuko 300 mcg prefilled syringe: 3 syringes/1 day
Granix 300 mcg single-dose vial: 4 vials/1 day	Releuko 480 mcg vial: 3 vials/1 day
Granix 480 mcg prefilled syringe: 3 syringes/1 day	Releuko 480 mcg prefilled syringe: 3 syringes/1 day
Granix 480 mcg single-dose vial: 3 vials/1 day	Rolvedon 13.2 mg prefilled syringe: 1 syringe/14 days
Leukine 250 mcg vial: 28 vials/14 days	Ryzneuta 20 mg prefilled syringe: 1 syringe/14 days
Neulasta 6 mg prefilled syringe: 1 syringe/14 days	Stimufend 6 mg prefilled syringe: 1 syringe/14 days
Neulasta 6 mg prefilled syringe kit: 1 kit/14 days	Udenyca 6 mg prefilled syringe: 1 syringe/14 days
Neupogen 300 mcg vial: 3 vials/1 day	Udenyca 6 mg auto-injector: 1 injection/14 days
Neupogen 300 mcg SingleJect: 3 syringes/1 day	Udenyca 6 mg onbody (syringe, with wearable injector): 1 syringe/14 days
Neupogen 480 mcg vial: 3 vials/1 day	Zarxio 300 mcg prefilled syringe: 3 syringes/1 day
Nivestym 300 mcg prefilled syringe: 3 syringes/1 day	Zarxio 480 mcg prefilled syringe: 3 syringes/1 day
Nivestym 480 mcg vial: 3 vials/1 day	Ziextenzo 6 mg prefilled syringe: 1 syringe/14 days
Nivestym 480 mcg prefilled syringe: 3 syringes/1 day	

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

PROVIDER PLEASE NOTE: SUBMISSION OF APPLICABLE DOCUMENTATION IS NECESSARY (I.E. CHART NOTES, DISEASE HISTORY, CURRENT/PAST THERAPY RECORD, COMPLETE BLOOD COUNT OR OTHER LABORATORY RESULTS) FOR COMPLETION OF REQUEST

Short-acting Granulocyte Colony-Stimulating Factors (G-CSFs)

PLEASE SELECT ONE OF THE FOLLOWING INDICATIONS FOR USE:

- Medication will be used as primary prevention of febrile neutropenia in members with non-myeloid malignancy meeting **ONE** of the following [**Length of authorization = 6 months**]:
 - Member is undergoing myelosuppressive chemotherapy with an expected incidence of febrile neutropenia greater than 20%
 - Member is undergoing myelosuppressive chemotherapy with an expected incidence of febrile neutropenia of 10% to < 20% **AND one or more** of the following co-morbidities (select all that apply):
 - Age >65 years receiving full dose intensity chemotherapy
 - Extensive prior exposure to chemotherapy
 - Previous exposure of pelvis, or other areas of large amounts of bone marrow, to radiation
 - Persistent neutropenia (ANC ≤ 1000/mm³)
 - Bone marrow involvement by tumor

(Continued on next page)

- ❑ Member has a condition that can potentially increase the risk of serious infection (i.e., HIV/AIDS with low CD4 counts)
- ❑ Recent surgery and/or open wounds
- ❑ Poor performance status
- ❑ Renal dysfunction (creatinine clearance <50 mL/min)
- ❑ Liver dysfunction (elevated bilirubin >2.0 mg/dL)
- ❑ Chronic immunosuppression in the post-transplant setting, including organ transplant

OR

- ❑ Member is 18 years of age or older, has a diagnosis of acute myeloid leukemia, **AND** filgrastim therapy is needed shortly following completion of induction or consolidation chemotherapy [**Length of authorization = 6 months**]

OR

- ❑ Member has been acutely exposed to myelosuppressive doses of radiation (Hematopoietic Acute Radiation Syndrome [H-ARS]) [**Length of authorization = Date of service only**]

OR

- ❑ Member has been diagnosed with a non-myeloid malignancy, **AND** will be receiving myeloablative chemotherapy following a bone marrow transplant [**Length of authorization = Date of service only**]

OR

- ❑ Medication will be used in apheresis collection of autologous hematopoietic progenitor cells [**Length of authorization = Date of service only**]

OR

- ❑ Member has been diagnosed with congenital, cyclic, or idiopathic neutropenia, **AND** is currently showing symptoms and incidence of complications (e.g., fever, infections, oropharyngeal ulcers) [**Length of authorization = 12 months**]

OR

- ❑ Treatment with filgrastim is needed as adjunctive treatment of febrile neutropenia when primary prophylaxis with a long-acting granulocyte colony stimulating factor is not given [**Length of authorization = 6 months**]

OR

- ❑ Adjunctive treatment of febrile neutropenia is considered clinically appropriate when at least **ONE** of the following risk factors are present (in the absence of prior growth factor use within the same chemotherapy cycle of treatment) (**select all that apply**) [**Length of authorization = 6 months**]:
 - ❑ Age > 65 years
 - ❑ Neutrophil recovery is expected to be delayed (greater than 10 days)
 - ❑ Neutropenia is profound (less than 0.1×10^9)
 - ❑ Active pneumonia
 - ❑ Sepsis syndrome (hypotension and/or multi-organ damage/dysfunction noted)

(Continued on next page)

- ❑ Invasive fungal or opportunistic infection
- ❑ Onset of fever during inpatient stay

NOTE: Febrile neutropenia is defined as an oral temperature > 38.3°C (101.0°F) or 2 consecutive readings of 38.0°C (100.4°F) for 1 hour, with an absolute neutrophil count less than 500 cells/μL (0.5 x 10⁹/L) or less than 1000 cells/μL and expected to fall below 500 cells/μL over the next 48 hours.

OR

- ❑ Member has a diagnosis of primary myelodysplastic syndrome, **AND** filgrastim therapy will be used in combination with epoetin to treat anemia [**Length of authorization = 6 months**]

OR

- ❑ Member has a diagnosis of non-Hodgkin lymphoma or multiple myeloma, **AND** filgrastim therapy will be used in combination with plerixafor for the collection of progenitor cells leading to subsequent autologous transplantation. [**Length of authorization = Date of service only**]

NOTE: Mozobil (plerixafor) requires prior authorization

❑ Granulocyte-macrophage Colony-Stimulating Factor (GM-CSF) [Leukine]

PLEASE SELECT ONE OF THE FOLLOWING INDICATIONS FOR USE:

- ❑ Member is 55 years of age or older, has a diagnosis of acute myeloid leukemia, **AND** sargramostim therapy is needed shortly after the completion of induction or repeat induction of chemotherapy [**Length of authorization = 6 months**]

OR

- ❑ Member is 2 years of age or older, **AND** sargramostim therapy is needed for faster reconstitution of myeloid to prepare for allogeneic bone marrow transplant (**NOTE: confirmation of HLA-matched donor status is required**) [**Length of authorization = 6 months**]

OR

- ❑ Member is 2 years of age or older, has undergone bone marrow transplant (allogeneic or autologous), **AND** sargramostim therapy is needed because there is delayed or failed neutrophil recovery [**Length of authorization = 6 months**]

OR

- ❑ Medication will be used in apheresis collection of autologous hematopoietic progenitor cells [**Length of authorization = Date of service only**]

OR

- ❑ Member is 2 years of age or older, has a diagnosis of acute lymphoblastic leukemia (ALL), Hodgkin lymphoma (HL), or non-Hodgkin lymphoma (NHL), **AND** sargramostim therapy is needed for faster reconstitution of myeloid following an autologous peripheral blood progenitor cell transplant or bone marrow transplant [**Length of authorization = 6 months**]

OR

(Continued on next page)

OR

- Member has been acutely exposed to myelosuppressive doses of radiation (Hematopoietic Acute Radiation Syndrome [H-ARS]) **[Length of authorization = Date of service only]**

OR

- Member has a diagnosis of high-risk neuroblastoma, **AND** sargramostim is needed for combination therapy with a with GD2-binding monoclonal antibody (i.e., dinutiximab or naxitamab) **[Length of authorization = 6 months]**

Long-acting Granulocyte Colony-Stimulating Factors (G-CSFs)

PLEASE SELECT ONE OF THE FOLLOWING INDICATIONS FOR USE:

- Medication will be used as primary prevention of febrile neutropenia in members with non-myeloid malignancy meeting **ONE** of the following **[Length of authorization = 6 months]**:
 - Member is undergoing myelosuppressive chemotherapy with an expected incidence of febrile neutropenia greater than 20%
 - Member is undergoing myelosuppressive chemotherapy with an expected incidence of febrile neutropenia of 10% to < 20% **AND one or more** of the following co-morbidities (select all that apply):
 - Age >65 years receiving full dose intensity chemotherapy
 - Extensive prior exposure to chemotherapy
 - Previous exposure of pelvis, or other areas of large amounts of bone marrow, to radiation
 - Previous/persistent neutropenia (ANC \leq 1000/mm³)
 - Bone marrow involvement by tumor
 - Patient has a condition that can potentially increase the risk of serious infection (i.e., HIV/AIDS with low CD4 counts)
 - Recent surgery and/or open wounds
 - Poor performance status
 - Renal dysfunction (creatinine clearance <50 mL/min)
 - Liver dysfunction (elevated bilirubin >2.0 mg/dL)
 - Chronic immunosuppression in the post-transplant setting, including organ transplant

OR

- Member has been acutely exposed to myelosuppressive doses of radiation (Hematopoietic Acute Radiation Syndrome [H-ARS]) **[Length of authorization = Date of service only]**

OR

- Medication will be used as secondary prevention of febrile neutropenia in members with non-myeloid malignancy, **AND** having experienced a neutropenic complication from a prior cycle of the same chemotherapy **[Length of authorization = 6 months]**

OR

- ❑ Treatment with requested medication is needed as adjunctive treatment of febrile neutropenia when primary prophylaxis is not given **[Length of authorization = 6 months]**

OR

- ❑ Adjunctive treatment of febrile neutropenia is considered clinically appropriate when at least **ONE** of the following risk factors are present (in the absence of prior growth factor use within the same chemotherapy cycle of treatment) (select all that apply) **[Length of authorization = 6 months]**:
 - ❑ Age > 65 years
 - ❑ Neutrophil recovery is expected to be delayed (greater than 10 days)
 - ❑ Neutropenia is profound (less than 0.1×10^9)
 - ❑ Active pneumonia
 - ❑ Sepsis syndrome (hypotension and/or multi-organ damage/dysfunction noted)
 - ❑ Invasive fungal or opportunistic infection
 - ❑ Onset of fever during inpatient stay

NOTE: Febrile neutropenia is defined as an oral temperature > 38.3°C (101.0°F) or 2 consecutive readings of 38.0°C (100.4°F) for 1 hour, with an absolute neutrophil count less than 500 cells/μL ($0.5 \times 10^9/L$) or less than 1000 cells/μL and expected to fall below 500 cells/μL over the next 48 hours

OR

- ❑ Treatment with requested medication is needed after bone marrow transplantation (BMT) failure or engraftment delay **[Length of authorization = 6 months]**

OR

- ❑ Medication will be used in apheresis collection of autologous hematopoietic progenitor cells **[Length of authorization = Date of service only]**

❑ **For medical necessity on a treatment purpose not listed, please provide clinical rationale and submit any chart notes/literature you feel would be pertinent in support of medical necessity:**

Medication being provided by Specialty Pharmacy – Proprium Rx

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****
****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****