AvMed

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions:</u> The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-877-535-1391</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If information provided is not complete, correct, or legible, authorization can be delayed</u>.

For Medicare Members: Medicare Coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: https://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx. Additional indications may be covered at the discretion of the health plan.

Macular Degeneration Drugs (Medical)

PREFERRED

Drug Requested: Check box below that applies.

□ Avastin[®] (bevacizumab) (J9035)

			<u> </u>		
	bevacizumab 1.25 mg/0.0)5 r	nL (3 mg/0.12 mL) intravi	itreal	injection (J9035)
			NON-PREFERRRED		
	Beovu® (brolucizumab) (J0179)		Byooviz [™] (ranibizumabnuna) (Q5124)		Cimerli [™] (ranibizumab-eqrn) (Q5128)
	Eylea® (aflibercept) (J0178)		Eylea® HD (aflibercept) (J0177)		Lucentis® (ranibizumab) (J2778)
	Pavblu [™] (aflibercept-ayyh) (Q5147)		Susvimo® (ranibizumab) (J2779)		Vabysmo®(faricimab-svoa) (J2777)
			NFORMATION: Authoriz	ation 1	may be delayed if incomplete.
	mber Name: mber AvMed #:			_ D:	ate of Birth:
Pre	scriber Name:				
					Date:
Off	ice Contact Name:				
Pho	ne Number:		Fax N	Numb	er:
NPl	[#:				

DRUG INFORMATION: Authorization may be delayed if incomplete.				
Diagnosis: ICD Code, if applicable:				
☐ Standard Review. In checking this box, the timeframe do or the member's ability to regain maximum function and	5 1			
□ Left Eye □ Right Eye	□ Both Eyes			
Preparations & Billable Units:				
Medication	Billable Units			
Beovu® (brolucizumab) 6 mg/0.05 mL solution	1 syringe = 6 billable units			
Byooviz [™] (ranibizumab-nuna) 0.5 mg/0.05 mL solution	1 vial = 5 billable units			
Cimerli® (ranibizumab-eqrn) 0.3 mg/0.05 mL solution	1 vial = 3 billable units			
Cimerli® (ranibizumab-eqrn) 0.5 mg/0.05 mL solution	1 vial = 5 billable units			
Eylea® (aflibercept) 2 mg/0.05 mL solution	1 syringe = 2 billable units			
Eylea® HD (afliberept) 8 mg/0.07 mL solution	1 vial = 8 billable units			
Lucentis® (ranibizumab) 0.3 mg/0.05 mL solution	1 syringe = 3 billable units			
Lucentis® (ranibizumab) 0.5 mg/0.05 mL solution	1 syringe = 5 billable units			
Susvimo® (ranibizumab) 10 mg/0.1 mL implant	1 vial/kit = 100 billable units			
Vabysmo®(faricimab-svoa) 6 mg/0.05 mL solution	1 vial = 60 billable units			
CLINICAL CRITERIA: Check below all that apply. support each line checked, all documentation, including lab provided or request may be denied. Avastin®/bevacizumab 1.25 mg/0.05 mL (3 m below all that apply. All criteria must be met for approve documentation, including lab results, diagnostics, and/or denied.	g/0.12 mL) intravitreal injection. Check al. To support each line checked, all			
☐ Provider has submitted member's baseline best correction	cted visual acuity (BCVA) score:			
 □ Member has been diagnosed with ONE of the follow □ Diabetic macular edema (DME) □ Diabetic retinopathy (DR) □ Neovascular (wet) age-related macular degenerate □ Macular edema following retinal vein occlusion (□ Myopic choroidal neovascularization (mCNV) 	ion (AMD)			
□ Neovascular glaucoma				

	Ot	her rare causes of choroidal neovascularization for ONE or more of the following conditions:
		Angioid streaks
		Choroiditis (including, but not limited to histoplasmosis induced choroiditis)
		Degenerative idiopathic myopia
		Retinal dystrophies
		Trauma
		Pseudoxanthoma elasticum
		Retinopathy of prematurity
		Other:
aj	pro	entis [®] , Byooviz [™] or Cimerli [™] . Check below all that apply. All criteria must be met for eval. To support each line checked, all documentation, including lab results, diagnostics, and/or notes, must be provided or request may be denied.
<u>Initi</u>	al A	Authorization: 12 months
	W	nich of the following medications is being requested for initial authorization?
		Lucentis®
		Byooviz [™]
		Cimerli™
	Pro	ovider has submitted member's baseline best corrected visual acuity (BCVA) score:
	Me	ember tried and failed at least 30 days of therapy with Avastin® or bevacizumab
	Pro	ovider has submitted chart notes to document treatment failure with the PREFERRED drug
	Me	ember has been diagnosed with ONE of the following labeled indications:
		Lucentis & Cimerli only - Diabetic macular edema (DME):
		☐ Intravitreal Dosing: 0.3 mg once a month
		Lucentis & Cimerli only - Diabetic retinopathy (DR):
		☐ Intravitreal Dosing: 0.3 mg once a month
		Neovascular (wet) age-related macular degeneration (AMD):
		☐ Intravitreal Dosing: 0.5 mg once a month
		Macular edema following retinal vein occlusion (MEfRVO):
		☐ Intravitreal Dosing: 0.5 mg once a month
		Myopic choroidal neovascularization (mCNV):
		☐ Intravitreal Dosing: 0.5 mg once a month for up to 3 months; may re-treat if necessary

a	ppro	entis [®] , Byooviz [™] or Cimerli [™] . Check below all that apply. All criteria must be met for val. To support each line checked, all documentation, including lab results, diagnostics, and/or notes, must be provided or request may be denied.
		orization: based on disease activity assessment
	-	nich of the following medications is being requested for reauthorization? Lucentis® Byooviz™ Cimerli™
	Pro	ovider has submitted member's BCVA score measured within the last 30 days:
1i	ne c	a [®] , Pavblu [™] . Check below all that apply. All criteria must be met for approval. To support each hecked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided uest may be denied.
<u>Initi</u>	al A	Authorization: 12 months
_		Eylea® Pavblu [™] Pavblu [™] Povider has submitted member's baseline best corrected visual acuity (BCVA) score:
		ember tried and failed at least 30 days of therapy with Avastin® or bevacizumab
		ovider has submitted chart notes to document treatment failure with the PREFERRED drug
		ember has been diagnosed with ONE of the following labeled indications: Neovascular (wet) age-related macular degeneration (AMD): Intravitreal Dosing: 2 mg (0.05 mL) once every 4 weeks for the first 12 weeks, followed by 2 mg (0.05 mL) once every 8 weeks Diabetic macular edema (DME): Intravitreal Dosing: 2 mg (0.05 mL) once every 4 weeks for the first 5 injections, followed by 2 mg (0.05 mL) once every 8 weeks Diabetic retinopathy (DR) with and/or without DME: Baseline Diabetic Retinopathy Disease Severity Scale (DRSS) Level: Intravitreal Dosing: 2 mg (0.05 mL) once every 4 weeks for the first 5 injections, followed by 2 mg (0.05 mL) once every 8 weeks Macular edema following retinal vein occlusion (MEfRVO):
	J	☐ Intravitreal Dosing: 2 mg (0.05 mL) once every 4 weeks

(Continued from previous page)

1i	ne c	a®, Pavblu™. Check below all that apply. All criteria must be met for approval. To support each hecked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided juest may be denied.
Rea	<u>uth</u>	orization: based on disease activity assessment
		hich of the following medications is being requested for initial authorization? Eylea® Pavblu [™]
		r diagnoses of Neovascular (wet) age-related macular degeneration (AMD) or Diabetic macular ema (DME):
		Provider has submitted member's BCVA score measured within the last 30 days:
		If no change in BCVA from baseline:
		☐ Maintenance Dose Intravitreal: 2 mg (0.05 mL) once every 8 weeks
		OR
		If increase in BCVA or increase presence of intraretinal or sub- retinal fluid or progression of pigment epithelial detachment):
		☐ Maintenance Dose Intravitreal: 2 mg (0.05 mL) once every 4 weeks
	Fo	r diagnosis of Diabetic retinopathy (DR) with and/or without DME:
		Provider has submitted member's Diabetic Retinopathy Disease Severity Scale (DRSS) Level recorded within the last 30 days:
		If DRSS level has decreased from baseline or member's baseline DRSS level was 10:
		☐ Maintenance Dose Intravitreal: Intravitreal Dosing: 2 mg (0.05 mL) once every 8 weeks
		OR
		If DRSS level has increased from baseline or no change has been observed:
		☐ Member does <u>NOT</u> have level 10 Disease Severity
		☐ Maintenance Dose Intravitreal: Intravitreal Dosing: 2 mg (0.05 mL) once every 4 weeks
cl	heck	a [®] HD. Check below all that apply. All criteria must be met for approval. To support each line ted, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or st may be denied.
<u>Initi</u>	al A	Authorization: 12 months
	Pro	ovider has submitted member's baseline best corrected visual acuity (BCVA) score:
	Me	ember tried and failed at least 30 days of therapy with Avastin [®] or bevacizumab AND Eylea [®]
		ovider has submitted chart notes to document treatment failure with the PREFERRED drug

(Continued from previous page)

	Me	emb	er h	as been diagnosed with ONE of the following labeled indications:
		Ne	ova	scular (wet) age-related macular degeneration (AMD):
				ravitreal Dosing: 8 mg once every 4 weeks for the first 3 doses, followed by ONE of the lowing (select requested dosing):
				8 mg once every 8 weeks
				8 mg once every 16 weeks
				Off-label dose: 8 mg every 4 weeks for 12 doses (Provider please note: if this dose is selected, it will NOT be approved, please prescribe another medication that is FDA approved for the requested indication)
		Di	abet	tic macular edema (DME):
				ravitreal Dosing: 8 mg once every 4 weeks for the first 3 doses, followed by ONE of the lowing (select requested dosing):
				8 mg once every 8 weeks
				8 mg once every 16 weeks
				Off-label dose: 8 mg every 4 weeks for 12 doses (Provider please note: if this dose is selected, it will NOT be approved, please prescribe another medication that is FDA approved for the requested indication)
		Di	abet	tic retinopathy (DR) with and/or without DME:
			Ba	seline Diabetic Retinopathy Disease Severity Scale (DRSS) Level:
				ravitreal Dosing: 8 mg once every 4 weeks for the first 3 doses, followed by ONE of the lowing (select requested dosing):
				8 mg once every 8 weeks
				8 mg once every 16 weeks
				Off-label dose: 8 mg every 4 weeks for 12 doses (Provider please note: if this dose is selected, it will NOT be approved, please prescribe another medication that is FDA approved for the requested indication)
cl	ieck	ed,	all (• Check below all that apply. All criteria must be met for approval. To support each line documentation, including lab results, diagnostics, and/or chart notes, must be provided or be denied.
ea	uth	<u>ori</u>	zat	ion: based on disease activity assessment.
	Fo	r dia	agno	oses of Neovascular (wet) age-related macular degeneration (AMD)
		Pro	ovid	ler has submitted member's BCVA score measured within the last 30 days:
		Se	lect	ONE of the following doses based on submission of member's BCVA score:
			If 1	no change in BCVA from baseline, maintenance dose intravitreal: 8 mg once every 8 weeks
			If	BCVA has improved from baseline, maintenance dose intravitreal: 8 mg once every 16 weeks
	Fo	r dia	agno	oses of Diabetic macular edema (DME)
		Pro	ovid	ler has submitted member's BCVA score measured within the last 30 days:
		Se	lect	ONE of the following doses based on submission of member's BCVA score:
			If	no change in BCVA from baseline, maintenance dose intravitreal: 8 mg once every 8 weeks
			If I	BCVA has improved from baseline: maintenance dose intravitreal: 8 mg once every 16 weeks

□ For diagnosis of Diabetic retinopathy (DR) with and/or without DME:
□ Provider has submitted member's Diabetic Retinopathy Disease Severity Scale (DRSS) Level recorded within the last 30 days:
☐ Select ONE of the following doses based on submission of member's DRSS level
☐ If DRSS level has decreased from baseline or member's baseline DRSS level was 10, maintenance dose intravitreal: 8 mg once every 12 weeks
☐ If DRSS level has increased from baseline or no change has been observed, maintenance dose intravitreal: 8 mg every 8 weeks
Beovu [®] . Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.
Initial Authorization: 3 months
☐ Provider has submitted member's baseline best corrected visual acuity (BCVA) score:
☐ Member tried and failed at least 30 days of therapy with Avastin® or bevacizumab
☐ Provider has submitted chart notes to document treatment failure with the PREFERRED drug
☐ Member has been diagnosed with ONE of the following labeled indications:
□ Neovascular (wet) age-related macular degeneration (AMD)
☐ Member has a diagnosis of Diabetic macular edema (DME)
☐ First Approval: Initial Dose Intravitreal: 6 mg once per month for 3 months
Beovu [®] . Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.
Reauthorization: based on disease activity assessment
☐ Provider has submitted member's BCVA score measured within the last 30 days:
☐ Member must meet ONE of the following:
☐ Disease activity is present (defined as loss of < 5 letters in BCVA score):
☐ Maintenance Dose Intravitreal: 6 mg once every 8 weeks
□ No disease activity is present:
☐ Maintenance Dose Intravitreal: 6 mg once every 12 weeks
□ Susvimo [™] . Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.
Initial Authorization: 12 months
☐ Provider has submitted member's baseline best corrected visual acuity (BCVA) score:

	Member is 18 years of age or older
	Member does NOT have ocular or periocular infection or active intraocular inflammation or conjunctival scarring
	Susvimo [™] will NOT be used with other ophthalmic VEGF inhibitors (unless supplemental treatment was approved
	Member tried and failed at least 30 days of therapy with Avastin® or bevacizumab
	Provider has submitted chart notes to document treatment failure with the PREFERRED drug
	Member tried and failed at least ONE of the following: □ Eylea® □ Beovu® □ Lucentis® □ Vabysmo®
	Member has a diagnosis of Neovascular (wet) age-related macular degeneration (AMD)
	Member has experienced disease stability or improvement following at least 2 injections in the same eye of either Beovu [®] , Eylea [®] , or Lucentis [®] prior to Susvimo [™] therapy
	Supplemental treatment to Susvimo [™] is allowed with Lucentis [®] only if ONE of the following are met: □ Decrease in visual acuity by half from the baseline visual acuity □ Increase of 150 µm or more in retinal thickness
cł	hecked , all documentation, including lab results, diagnostics, and/or chart notes, must be provided or equest may be denied.
eat	uthorization: 12 months (based on disease activity assessment)
	Medication has <u>NOT</u> caused toxicity to the eye (e.g., endophthalmitis, rhegmatogenous retinal detachment, implant dislocation, vitreous hemorrhage, conjunctival erosion, conjunctival retraction, and conjunctival blebs)
	Member has experienced a beneficial response to therapy (e.g., improvement in the baseline best corrected visual acuity (BCVA), and does not show loss of more than 20 letters in a BCVA (best corrected visual acuity)
cł	abysmo [®] . Check below all that apply. All criteria must be met for approval. To support each line hecked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or equest may be denied.
<u>iiti</u>	al Authorization: 6 months.
	Provider has submitted member's baseline best corrected visual acuity (BCVA) score:
	Member tried and failed at least 30 days of therapy with Avastin® or bevacizumab (submit chart notes to document treatment failure)

	Member has been diagnosed with ONE of the following labeled indications:
	□ Neovascular (wet) age-related macular degeneration (AMD):
	☐ Intravitreal Dosing: 6 mg once every 4 weeks for 4 doses, followed by ONE of the following
	dosing regimens:
	□ Every 16 weeks
	□ Every 12 weeks
	□ Every 8 weeks
	□ Diabetic macular edema (DME):
	☐ Intravitreal Dosing: 6 mg once every 4 weeks for 6 doses, followed by 6 mg once every 8 weeks
	Therapy will NOT be used with other ophthalmic VEGF inhibitors (e.g., aflibercept, brolucizumab-dbll,
_	ranibizumab, pegaptanib, bevacizumab)
ch	abysmo [®] . Check below all that apply. All criteria must be met for approval. To support each line necked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or quest may be denied.
	· · ·
	y Reauthorization: 3 months. Applicable for patients with an insufficient response
	ng initial therapy administered every 4 weeks for at least 4 doses requesting
conti	inuation of every 4-week dosing.
	Provider has submitted progress notes which document patient has experienced an insufficient response every 4-week dosing as detected by clinical exam, optical coherence tomography or decrease in best corrected visual acuity score
ch	abysmo [®] . Check below all that apply. All criteria must be met for approval. To support each line necked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or quest may be denied.
Reau	uthorization: 12 months (based on disease activity assessment).
Prov	vider Please Note: Patients with loss of response to maintenance therapy
adm	inistered at less frequent intervals may increase the dosing frequency in a stepwise
man	ner until response is regained.
	Medication has <u>NOT</u> caused toxicity to the eye (e.g., endophthalmitis, rhegmatogenous retinal detachment, implant dislocation, vitreous hemorrhage, conjunctival erosion, conjunctival retraction, and conjunctival blebs)
	Member has experienced a beneficial response to therapy (e.g., resolution of edema based on the central subfield thickness (CST) of the macula as measured by optical coherence tomography is achieved, improvement in the baseline best corrected visual acuity (BCVA)
	Provider will administer requested medication via ONE of the following dosing regimens:
	□ Every 16 weeks
	□ Every 12 weeks
	□ Every 8 weeks
	(Continued on next page)
	(Constitute on none puge)

	Location/site of drug administration:
	NPI or DEA # of administering location:
	OR
	Specialty Pharmacy
eviev eatm	gent reviews: Practitioner should call AvMed Pre-Authorization Department if they believe a standard would subject the member to adverse health consequences. AvMed's definition of urgent is a lack of the nent that could seriously jeopardize the life or health of the member or the member's ability to regain num function.
	*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.** evious therapies will be verified through pharmacy paid claims or submitted chart note.