AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-305-671-0200. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process can be delayed.

Drug Requested: Benlysta® (belimumab) Subcutaneous Injection (Pharmacy)

MEMBER & PRI	ESCRIBER INFORMATION:	Authorization may be delayed if incomplete.	
Member Name:			
Member Sentara #:			
Prescriber Name:			
Prescriber Signature:			
Office Contact Name:			
Phone Number:			
NPI #:			
DRUG INFORMA	ATION: Authorization may be dela	yed if incomplete.	
Drug Name/Form/Str	ength:		
Dosing Schedule:		Length of Therapy:	
Diagnosis:	nosis: ICD Code, if applicable:		
Weight (if applicable):		Date weight obtained:	
Recommended Dos	sing:		
	A J-14- (A4	Dadiatria Datianta 5 to loss than 10	

Diagnosis	Adults (Auto-injector or Prefilled syringe)	Pediatric Patients 5 to less than 18 years of age (Auto-injector only)	
SLE	200 mg once weekly	 Patients ≥ 40 kg: 200 mg once weekly Patients 15 kg to <40 kg: 200 mg once every 2 weeks 	
Lupus Nephritis	400 mg once weekly x 4 doses, followed by 200 mg once weekly	Safety and efficacy of subcutaneous administration has not been established	

Quantity Limits: 200 mg once weekly (4 injections per 28 days)

CLINICAL CRITERIA/DIAGNOSIS: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied. Check box below for the Diagnosis that applies.

	iagnosis - active systemic lu andard therapy	pus erythematosus (SLE) in pa	tients who are receiving		
	al Authorization: 12 month	18			
	Prescribed by or in consultation	with a rheumatologist			
	•	er is 5 years of age or older with a diagnosis of active, autoantibody-positive SLE confirmed by f the following (submit lab results):			
	☐ anti-nuclear antibody (ANA)	titer $\geq 1:80$			
	□ anti-double stranded DNA (a	$nti-dsDNA) \ge 30 \text{ IU/mL}$			
	Member's SLE activity has been confirmed by ONE of the following (submit results):				
	□ Safety of Estrogen in Lupus National Assessment – Systemic Lupus Erythematosus Disease Activity Index (SELENA-SLEDAI) score of 6-12				
	□ ≥2 British Isles Lupus Assess	sment Group (BILAG) B organ domair	1 scores		
		lowing and is established on two of the trotes documenting therapy trials v	<u> </u>		
	□ mycophenolate	□ hydroxychloroquine	□ azathioprine		
	□ cyclophosphamide	□ methotrexate	□ cyclosporine		
	□ corticosteroids	□ Other			
		the following limitations to therapy: se sis of progressive multifocal leukoence	•		
	iagnosis - active systemic lu andard therapy	pus erythematosus (SLE) in pa	tients who are receiving		
suppo		eck below all that apply. All criteria n tation, including lab results, diagnostic			
	All initial authorization criteria continues to be met				
	☐ Member's response to therapy has been confirmed by ONE of the following (submit results) :				
	•	National Assessment – Systemic Lupus core has improved by and/or maintaine	•		
	☐ No new BILAG-A organ don	nain score OR 2 new BILAG-B organ	domain scores		
		rable side effects such as serious infect phalopathy (PML), malignancy, severe infusion reactions	· · · · · ·		

□ D	iagnosis - active lupus nephritis in adults who are receiving standard therapy
<u>Initi</u>	al Authorization: 12 months
	Prescribed by or in consultation with a nephrologist or rheumatologist
	Member is 18 years of age or older with a diagnosis of active lupus nephritis Class III, IV, or V as confirmed by renal biopsy
	Member's diagnosis of active, autoantibody-positive SLE was confirmed by ONE of the following (submit lab results) :
	 □ anti-nuclear antibody (ANA) titer ≥ 1:80 □ anti-double stranded DNA (anti-dsDNA) ≥ 30 IU/mL
	Member has active renal disease and has received standard therapy for the last 90 days with corticosteroids along with <u>ONE</u> of the following (chart notes documenting established therapy must be submitted):
	□ mycophenolate□ cyclophosphamide
	Provider must obtain a baseline measurement of <u>ONE</u> of the following collected within the last 30 days (labs must be submitted):
	□ urine protein:creatinine ratio (uPCR)□ urine protein
	Member does <u>NOT</u> have any of the following limitations to therapy: severe active central nervous system lupus, current or previous diagnosis of progressive multifocal leukoencephalopathy (PML), or concurrent use with other biologics
ı Di	agnosis - active lupus nephritis in adults who are receiving standard therapy
suppo	thorization: 12 months. Check below all that apply. All criteria must be met for approval. To reach line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be led or request may be denied.
	All initial authorization criteria continues to be met
	Member has had improvement from baseline and/or stabilization since last approval of one of the following (submit current labs completed within the last 30 days): urine protein:creatinine ratio (uPCR)
	urine protein
	Member has an absence of intolerable side effects such as serious infections, signs or symptoms of progressive multifocal leukoencephalopathy (PML), malignancy, severe hypersensitivity reactions/anaphylaxis, or serious infusion reactions
Med	lication being provided by a Specialty Pharmacy – Proprium Rx

**Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *