AvMed

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-305-671-0200</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

<u>Drug Requested</u>: Enspryng[™] (satralizumab-mwge) (Pharmacy)

Neuromyelitis Optica Spectrum Disorder (NMOSD)

MEMBER & PRESCRIBER IN	NFORMATION: Authorization may be delayed if incomplete.			
Member Name:				
Member AvMed #:	Date of Birth:			
Prescriber Name:				
Prescriber Signature:	Date:			
Office Contact Name:				
Phone Number: Fax Number:				
DEA OR NPI #:				
DRUG INFORMATION: Author	orization may be delayed if incomplete.			
Drug Form/Strength:				
Dosing Schedule:	Length of Therapy:			
Diagnosis:	ICD Code, if applicable:			
Weight:	Date:			
	lose: 120 mg once every 2 weeks for 3 doses (weeks 0, 2, and 4), once every 4 weeks. Maximum quantity: 120mg every 4 weeks.			
each line checked, all documentation, in	below all that apply. All criteria must be met for approval. To support acluding lab results, diagnostics, and/or chart notes, must be provided or			
Initial Authorization Approval:	6 months			
☐ Prescribing physician must be a	neurologist			
AND				
☐ Member must be 18 years of age	or older			
AND				

		lust submit medical records (e.g. chart notes, laboratory values, etc.) to support a diagnosis of euromyelitis Optica Spectrum Disorder (NMOSD) confirming all of the following:				
	st medical history of one of the following:					
			Optic neuritis			
			Acute myelitis			
			Area postrema syndrome; episode of otherwise unexplained hiccups or nausea and vomiting			
			Acute brainstem syndrome			
			Symptomatic narcolepsy or acute diencephalic clinical syndrome with NMOSD-typical diencephalic MRI lesions			
			Symptomatic cerebral syndrome with NMOSD-typical brain lesions			
			AND			
			sitive serologic test for anti-aquaporin-4 immunoglobulin G (AQP4-IgG)/NMP-IgG antibodies ust submit lab results)			
			AND			
		Dia	agnosis of multiple sclerosis or other diagnoses have been ruled out			
			AND			
	Member has a history of at least one relapse during the previous 12 months prior to initiating Enspryng or member has a history of at least two relapses during the previous 24 months, at least one relapse occurring within the past 12 months prior to initiating Enspryng [™]					
	net	arol ona	torical relapse is defined as a new onset of neurologic symptoms or worsening of existing ogic symptoms with an objective change on neurologic examination (clinical findings, magnetic nee imaging findings, or both) that persist for more than 24 hours and/or the new onset of ogic symptoms or worsening of existing neurologic symptoms that require treatment.}			
			AND			
			er must have documentation of an inadequate response or intolerance with rituximab during the 12 s prior to initiating Enspryng [™]			
			OR			
			nber has a documented contraindication to rituximab, member must have failed the following ent during the 12 months prior to initiating Enspryng TM			
			ember must have failed at least 2 immunosuppressive therapies (e.g., azathioprine, cyclosporine, cophenolate, etc.)			
			OR			
			ember must have failed at least 1 immunosuppressive therapy and required chronic plasmapheresis plasma exchange (PE) or intravenous immunoglobulin (IVIG)			
			AND			
			(Continued on next page)			
			(Continued on next page)			

	Member does not have an active infection, including clinically important localized infections
	AND
	Provider has submitted a baseline liver transaminase and neutrophil count prior to treatment and will continue to monitor throughout treatment
	AND
	Member has been evaluated and screened for the presence of latent TB infection prior to initiating treatment
	AND
	Member has been evaluated and screened for the presence hepatitis B virus (HBV) prior to initiating treatment
	AND
	Enspryng [™] will not be used in combination with disease-modifying therapies for the treatment of multiple sclerosis (e.g. Gilenya (fingolimod), Tecfidera (dimethyl fumarate), Ocrevus (ocrelizumab), etc.)
	AND
	Enspryng [™] will not be used in combination with other complement inhibitor therapy (e.g.,eculizumab), IL6-inhibitors (e.g., toclizumab), anti-CD20-directed antibody therapy (e.g., rituximab) or anti-CD19 directed antibody therapy (ineblizumab-cdon)
prov	thorization Approval: 12 months. Check below all that apply. All criteria must be met for val. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, be provided or request may be denied.
	Member continues to meet the initial criteria
	AND
	Absence of unacceptable toxicity from therapy (i.e. tuberculosis (TB) infections, hepatitis B reactivation, infusion reactions, serious infections)
	AND
	Provider must submit clinical notes documenting clinical improvement (fewer relapses from baseline) or stabilization of patient relapses while on Enspryng [™] therapy

treat NMOSD or exacerbation of symptoms while on therapy will be considered as treatment failure.

(Continued on next page)

Note: Add on, dose escalation of immunosuppressive therapy, or additional rescue therapy from baseline to

PA Enspryng (AvN	(led)
(Continued from previous p	age)

3 6 3 4 4			~		TO 1	-		_
Medication	heing	nrovided b	V	necialty	Pharmacy	- Pro	nriiiml	٧v
MICHICALIUM		DI U IIUCU D	' T	Deciales	1 1141 11140	110	DIIMIII	

Not all drugs may be covered under every Plan

If a drug is non-formulary on a Plan, documentation of medical necessity will be required.

**Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.