

AvMed

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-877-535-1391**. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization can be delayed.**

For Medicare Members: Medicare Coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx>. Additional indications may be covered at the discretion of the health plan.

Drug Requested: Casgevy® (exagamglogene autotemcel) (J3392) (Medical)
β - Thalassemia

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member AvMed #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight (if applicable): _____ Date weight obtained: _____

Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

- Casgevy® is supplied in one or more vials (one carton contains a single lot consisting of 1 to 9 vials) containing a frozen suspension of genome edited autologous CD34+ cells in a cryo-preserved medium containing 5% DMSO and dextran 40 [NDC 51167-290-09]
- The minimum recommended dose of Casgevy® is 3×10^6 CD34+ cells per kg of body weight

B. Max Units (per dose and over time) [HCPCS Unit]:

- One treatment (dose) per lifetime

(Continued on next page)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Coverage will be provided for one treatment course and may NOT be renewed.

- Member is ≥ 12 years of age
- Medication is prescribed by a hematologist or a stem cell transplant physician
- Member has **NOT** received a gene therapy for beta-thalassemia in the past (**verified by medical paid claims**) [**NOTE:** If no claim for Casgevy[®] or Zynteglo[™] (betibeglogene autotemcel intravenous infusion) is present (or if claims history is not available), the prescribing physician confirms that the member has not previously received Casgevy[®] or Zynteglo[™]]
- According to the prescribing physician, a hematopoietic stem cell transplantation is appropriate for the member
- Member meets **ONE** of the following:
 - Member does **NOT** have a Human Leukocyte Antigen (HLA)-matched donor
 - Member has an HLA-matched donor, but the individual is not able or is not willing to donate
- Member has **ONE** of the following genotypes as confirmed by genetic testing (**submit documentation**):
 - Non- β^0/β^0 genotype [**NOTE:** Examples include β^0/β^+ , β^E/β^0 , and β^+/β^+]
 - β^0/β^0 genotypes [**NOTE:** Other examples include $\beta^0/\beta^{+(IVS-I-110)}$ and $\beta^{+(IVS-I-110)}/\beta^{+(IVS-I-110)}$]
- Member is transfusion-dependent, as defined by meeting **ONE** of the following (**submit documentation**):
 - Receipt of transfusions of ≥ 100 mL of packed red blood cells per kg of body weight per year in the previous 2 years
 - Receipt of transfusions of ≥ 10 units of packed red blood cells per year in the previous 2 years
- Member meets **BOTH** of the following:
 - Member has been evaluated for the presence of severe iron overload (**submit documentation**)
 - Member does **NOT** have evidence of severe iron overload [**NOTE:** Examples include abnormal myocardial iron results (a T2*-weighted magnetic resonance imaging measurement of myocardial iron of less than 10 msec); high liver iron concentration (≥ 15 mg/g); liver biopsy results suggest abnormalities; or clinical evidence of organ damage (e.g., endocrine comorbidities)]
- Member does **NOT** currently have an active bacterial, viral, fungal, or parasitic infection
- Member does **NOT** have the following:
 - Prior or current malignancy or myeloproliferative disorder or significant immunodeficiency disorder [**NOTE:** This does not include adequately treated cone biopsied in situ carcinoma of the cervix uteri and basal or squamous cell carcinoma of the skin]
 - Advanced liver disease (**submit documentation**) [**NOTE:** Examples include alanine transaminase or aspartate transaminase greater than three times upper limit of normal, direct bilirubin value greater than three times upper limit of normal, active hepatitis, extensive bridging fibrosis, or cirrhosis]

(Continued on next page)

- According to the prescribing physician, member will have been discontinued from iron chelation therapy for at least 7 days prior to myeloablative conditioning [**NOTE**: Examples of iron chelators used for this condition include deferoxamine injection, deferi-prone tablets or solution, and deferasirox tablets]
- According to the prescribing physician, member meets **ALL** the following:
 - Member will undergo mobilization, apheresis, and myeloablative conditioning
 - A hematopoietic stem cell mobilizer will be utilized for mobilization [**NOTE**: Filgrastim products are examples of a granulocyte-colony stimulating factor therapy and Mozobil[®] (plerixafor subcutaneous injection) is an example of a hematopoietic stem cell mobilizer]
 - Busulfan will be used for myeloablative conditioning
 - Total hemoglobin level is ≥ 11 g/dL at **BOTH** of the following timepoints:
 - Prior to mobilization
 - Prior to myeloablative conditioning
- Prior to collection of cells for manufacturing, member cellular screening is negative for **ALL** the following (**submit documentation**):
 - Human immunodeficiency virus-1 and -2
 - Hepatitis B virus
 - Hepatitis C virus
 - Human T-lymphotrophic virus-1 and -2
- According to the prescribing physician, member meets **ONE** of the following:
 - Member is a female of reproductive potential and meets **BOTH** of the following:
 - A negative serum pregnancy test will be confirmed prior to the start of mobilization and re-confirmed prior to myeloablative conditioning
 - Member will use an effective method of contraception from the start of mobilization through at least 6 months after administration of Casgevy[®]
 - Member is a male of reproductive potential and will use an effective method of contraception from the start of mobilization through at least 6 months after administration of Casgevy[®]
- Member's current body weight has been obtained within 30 days (**submit documentation**)

Medication being provided by: Please check applicable box below.

Location/site of drug administration: _____

NPI or DEA # of administering location: _____

OR

Specialty Pharmacy – Proprium Rx

(Continued on next page)

For urgent reviews: Practitioner should call AvMed Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. AvMed's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****